

# Aerobic reserve and physical functional performance in older adults

SCOTT W. ARNETT<sup>1</sup>, JENNIFER H. LAITY<sup>2</sup>, SUBODH K. AGRAWAL<sup>3</sup>, M. ELAINE CRESS<sup>4</sup>

<sup>1</sup>Department of Physical Education and Recreation, Western Kentucky University, Bowling Green, KY, USA

<sup>2</sup>Bay Sport Inc., San Jose, CA, USA

<sup>3</sup>Athens Heart Center, Athens, GA, USA

<sup>4</sup>Department of Kinesiology, Gerontology Center, Aging and Physical Performance Laboratory, University of Georgia, Athens, GA, USA

Address correspondence to: Scott W. Arnett. Fax: (+1) 270-745-6034. Email: scott.arnett@wku.edu

## Abstract

**Background:** older adults can be limited in their performance of daily tasks due to an inadequate aerobic capacity. Aerobic capacity below minimum physiological thresholds required to maintain independence leaves older adults with little, or no, aerobic reserve.

**Objective:** the aim of this study was to measure functional performance and aerobic reserve in older adults during the serial performance of daily tasks.

**Subjects:** twenty-nine ( $n = 29$ ) men and women ( $n = 23$  females) 70–92 years of age participated in this study.

**Methods:** performance based physical function was assessed using the Continuous-Scale Physical Functional Performance test (CS-PFP). A Cosmed K4b<sup>2</sup>™ portable metabolic system was used to measure  $VO_{2PEAK}$  and oxygen uptake during the serial performance of a battery of daily tasks ( $VO_{2PFP}$ ). Aerobic reserve was calculated as the difference between  $VO_{2PEAK}$  and  $VO_{2PFP}$ .

**Results:** the correlation coefficient between aerobic reserve and functional performance was  $r = 0.50$  ( $P = 0.006$ ). Participants utilized  $32.2 \pm 8.1\%$ ,  $42.7 \pm 10.8\%$ , and  $50.3 \pm 12.3\%$  of  $VO_{2PEAK}$  for the low, moderate, and high workloads of the CS-PFP, respectively.

**Conclusions:** light housework and carrying groceries require 40 to 50% of peak oxygen consumption. This information can be used by clinicians and health professionals working with older adults as a guide to how much aerobic fitness is needed to perform ADLs and maintain independence. These can then be used as guides for assessment and for setting training goals in older adults.

**Keywords:**  $VO_{2PEAK}$ , CS-PFP, threshold, daily tasks, elderly

Aerobic capacity between 18 and 20 ml/kg/min [1–3] has been defined as an aerobic threshold below which there is a reduced probability of living independently [1]. For each milliliter of oxygen uptake below the threshold, there is an associated 8-fold decline in physical function [1]. Once above the threshold, higher aerobic capacity provides an aerobic reserve, and thus a margin of safety for discretionary activities such as active leisure activities [4]. Older adults with little or no aerobic reserve may be perilously close to, or fall below, minimum physiological thresholds required to perform daily tasks [5].

It is more practical and sensible to measure the work of intermittent activities rather than single, isolated activities [6]. To sustain daily function, particularly by those with limited

aerobic capacity, tasks are punctuated by rest breaks [6]. The continuous-scale physical functional performance (CS-PFP) test mimicking the performance of functional tasks in daily life is associated with the probability of living independently [1]. Understanding the aerobic cost of serial performances of daily tasks mimicking real-life situations relative to aerobic capacity may provide insights into the importance of an individual's aerobic capacity and reserves for independent living.

The purpose of this article was to determine (i) the oxygen cost of physical function, (ii) aerobic reserve, and (iii) capacity of physical functional performance. We hypothesised that older adults with higher aerobic reserves will have a higher physical functional performance capacity.

## Methods

### Participants

Participants were recruited from local social groups, assisted living facilities and using public service announcements on the radio. Forty-four older men and women were recruited to participate in this study of which seven were not medically cleared and seven chose not to participate for personal reasons. One participant's data file was eliminated due to data corruption, resulting in a final sample of 29 participants (23 females), representing 66% of the original sample. Testing was conducted on three separate days with a minimum of two rest days between each day of testing. Exclusion criteria were evaluated by their personal physician based on the following criteria: uncontrolled cardiovascular disease; uncontrolled diabetes; vigorous, aerobic exercise more than 3 days/week, 30 min/session; recent bone fracture or joint replacement; diseases with a variable course (e.g. rheumatoid arthritis); severe osteopenia; cardiovascular intolerance to exercise; severe psychiatric illness; and severe hypertension. This study was approved by the Institutional Review Board of the University of Georgia and all participants provided written informed consent.

### Performance-based physical function

Physical function was assessed during the first visit using the CS-PFP test, which has established validity and reliability [7]. The CS-PFP protocol and scoring system have been published elsewhere [8] and on the Web at <http://www.coe.uga.edu/cs-pfp/>. A brief description is included here for the reader's convenience. The CS-PFP protocol includes a standard dialogue and environment for the administration of 16 daily life tasks, quantified by time, distance and weight. Participants were instructed to perform the test at maximal effort within the bounds of safety and comfort. Tasks reflective of the domains of upper body strength (UBS), upper body flexibility (UBF), lower body strength (LBS), balance and coordination (BALC), and endurance (END), as determined using a modified Delphi procedure [8], are scaled from 0 to 100 and averaged to provide domain scores. The CS-PFP yields a total summary score (CS-PFP total), which is the average of all variables [8]. The sum of time taken to complete all timed tasks was PFP-TIME. Performance data were scored using the Web-based data reduction program at <http://www.coe.uga.edu/cs-pfp/>. The CS-PFP tasks are separated into workload categories based on the difficulty of each task as shown in Table 3. On a separate visit, participants performed a modified version of the CS-PFP (CS-PFP<sub>MOD</sub>) in order to measure oxygen uptake while performing activities of daily living. The CS-PFP<sub>MOD</sub> protocol is described in the section Oxygen Cost of Standardised Load Physical Functional Performance (VO<sub>2PFP</sub>), below.

### Peak oxygen uptake (VO<sub>2PEAK</sub>)

During the second visit, VO<sub>2PEAK</sub> was measured with a Cosmed K4b<sup>2</sup> portable metabolic system (Cosmed

S.R.L., Rome, Italy) during a modified Balke continuous, incremental treadmill protocol [9]. The Cosmed K4b<sup>2</sup> portable metabolic system was used to collect continuous, breath-by-breath values for oxygen uptake through indirect calorimetry. At rest and over a range of exercise intensities, the Cosmed K4b<sup>2</sup> has been shown to provide similar values of oxygen uptake, carbon dioxide production, minute ventilation and respiratory exchange ratio (RER) as those obtained by the Douglas bag method or a metabolic unit [10, 11]. Calibration of the Cosmed K4b<sup>2</sup> was conducted prior to each test following the manufacturer's guidelines [12].

Participants were familiarised with the face mask and walking on the treadmill, separately, prior to the VO<sub>2PEAK</sub> test. To determine the pace for the test, participants were instructed to walk at their fastest, comfortable pace during a 2 min warm-up. The treadmill grade was increased 2–3% every 2 min while speed remained at the participants' fastest, comfortable pace (range = 1.2–4.0 mph, mean  $\pm$  SD = 2.5  $\pm$  0.7 mph). A 12-lead Quinton 4000 (Quinton 4000, Bothell, WA) electrocardiogram was monitored by a cardiologist, and heart rate, blood pressure and rating of perceived exertion (RPE) (Borg scale 6–20) were collected at the end of each 2 min stage. Test termination occurred when participants reached volitional fatigue or demonstrated symptoms of cardiovascular incompetence according to ACSM guidelines [13]. The highest 30 s average VO<sub>2</sub> achieved during the treadmill test was considered VO<sub>2PEAK</sub>. The value was determined by calculating the average of all breaths during the last 30 s period of a stage. Two of the following three criteria are often used to establish maximal effort: (i) a maximum heart rate within  $\pm$ 10 beats of age-predicted maximum heart rate [14], (ii) an respiratory exchange ratio (RER) value >1.0 [15], and/or (iii) a RPE of 18 or higher on the Borg 6–20 scale [14, 16]. VO<sub>2PEAK</sub> was used as none of the participants in this study demonstrated a plateau in oxygen uptake during the test. However, 69% ( $n$  = 20) achieved the maximal heart rate criterion, 55% ( $n$  = 16) achieved the RER criterion, and 79% ( $n$  = 23) achieved the RPE criterion with 93% ( $n$  = 27) of all participants meeting at least one criterion.

### Oxygen cost of standardised load physical functional performance (VO<sub>2PFP</sub>)

During the third visit, the Cosmed K4b<sup>2</sup> measured oxygen uptake during performance of the CS-PFP<sub>MOD</sub>. CS-PFP<sub>MOD</sub> is used to distinguish the administration of the modified version of the CS-PFP used during assessment of oxygen uptake during performance of functional tasks from the original CS-PFP [8]. To standardise the workload across subjects, a standardised weight was used for the three tasks labelled 'b' in Table 3. The weight was determined by taking the lowest quartile of weight carried by older adults in previous research [7]. A 3 min walk at the participant's usual pace was substituted for the 6 min walk as normal pace is

## Aerobic reserve and functional performance

the most efficient (lowest oxygen uptake), reproducible, and individualised [17]. Donning and removing a jacket was omitted because the Cosmed K4b<sup>2</sup> unit was worn on the torso. Collection and analyses of gas volumes and composition were carried out using the same procedures previously described for the VO<sub>2PEAK</sub> test.

### VO<sub>2PFP</sub> and aerobic reserve analysis

Erratic zero values within the VO<sub>2</sub> data were replaced with the average of the data point immediately before and after each zero, and a moving window average (window size = 11 data points) was used to smooth the VO<sub>2</sub> data. Group analysis of VO<sub>2</sub> data was performed utilising three methods: (i) VO<sub>2PFP</sub> was determined by averaging VO<sub>2</sub> from the first task until completion of the last task, including VO<sub>2</sub>, between tasks of the CS-PFP<sub>MOD</sub>; (ii) low, moderate, and high was determined by averaging VO<sub>2</sub> measured from the beginning of the first task for each workload to the beginning of the first task of the next workload of the CS-PFP<sub>MOD</sub>; and (iii) average VO<sub>2</sub> for each task was determined where VO<sub>2</sub> was averaged from the beginning of one task to the beginning of the next task of the CS-PFP<sub>MOD</sub>. VO<sub>2PFP</sub> and the average VO<sub>2</sub> of each workload (low, moderate and high) were calculated as a percentage of VO<sub>2PEAK</sub>. Aerobic reserve PFP (AR-PFP) was determined by subtracting VO<sub>2PFP</sub> from VO<sub>2PEAK</sub> (VO<sub>2PEAK</sub> — VO<sub>2PFP</sub>).

### Statistical analysis

Analyses were performed using SPSS for Windows (Version 14.0, SPSS Inc., Chicago, IL). Independent samples *t*-test were used to determine differences between males and females on selected characteristics and physiological and functional performance variables. Levene's and Bonferroni adjustments ( $\alpha = 0.005$ ) were used when necessary. VO<sub>2PEAK</sub>, VO<sub>2PFP</sub> and CS-PFP total data were evaluated for normality. Pearson's product moment correlation was calculated to evaluate the association between age and VO<sub>2PEAK</sub> and CS-PFP total and the association of AR-PFP and the following variables: END, CS-PFP total, and PFP-TIME. A point biserial correlation coefficient ( $r_{pb}$ ) was calculated to evaluate the relationships between living status and the variables mentioned previously. An  $\alpha$  level of 0.05 was used to denote significance on correlation analyses.

### Results

Selected characteristics of the participants are reported in Table 1. The age of the participants in this study ranged from 70 to 92 years. Females represented 79% ( $n = 23$ ) of the sample, and 17.2% were residents of a retirement community. The sample was generally free of prescribed medications (76.6%) or on one medication (23.4%). CS-PFP total and domain scores are reported in Table 1. Except for height (men = 175.1 ± 9.2 cm; women = 160.5 ± 7.5 cm;  $P < 0.001$ ), data from men ( $n = 6$ ) and women ( $n = 23$ )

**Table 1.** Means ± SD of selected characteristics and continuous-scale physical functional performance characteristics

	Mean ± SD
Age (years)	76.2 ± 6.2
Height (cm)	163.5 ± 9.8
Weight (kg)	70.8 ± 10.8
BMI	26.55 ± 4.1
Independent	82.8%
CS-PFP total	52.1 ± 12.7
UBS	60.8 ± 11.1
LBS	44.6 ± 12.8
UBF	70.1 ± 11.9
BALC	47.5 ± 15.0
END	53.1 ± 14.3

SD, standard deviation; independent, detached single-family dwelling or apartment; CS-PFP total, continuous-scale physical functional performance total score; UBS, upper body strength; LBS, lower body strength; UBF, upper body flexibility; BALC, balance/coordination; END, endurance; CS-PFP total, UBS, LBS, UBF, BALC, and END are scored out of a total 100 possible points.

were not statistically different. All means, standard deviations and correlations reported include data from both males and females.

The data appeared to have normal distributions for VO<sub>2PEAK</sub>, VO<sub>2PFP</sub>, and CS-PFP. Selected physiologic characteristics of the group are listed in Table 2. Average oxygen uptake and the percent of VO<sub>2PEAK</sub> increased with the workload level of the CS-PFP<sub>MOD</sub>. The oxygen uptakes of individual tasks performed during the CS-PFP<sub>MOD</sub> protocol are presented in Table 3.

Correlations between AR-PFP and END, CS-PFP total, and PFP-TIME were  $r = 0.45$  ( $P < 0.05$ ),  $r = 0.50$  ( $P < 0.01$ ), and  $r = -0.43$  ( $P < 0.05$ ), respectively. The correlation between living status (living status -1 = assisted living, 2 = community dweller) and CS-PFP total was  $r_{pb} = 0.52$  ( $P < 0.01$ ). Age was significantly related to both VO<sub>2PEAK</sub> and CS-PFP total,  $r = -0.62$  and  $-0.50$ , respectively.

**Table 2.** Means ± SD of selected physiologic characteristics

( $n = 29$ )	VO <sub>2</sub> (ml/kg/min)	Percent of VO <sub>2PEAK</sub> (%)
VO <sub>2PEAK</sub>	19.8 ± 6.3	
VO <sub>2PFP</sub>	7.5 ± 1.4	40.3 ± 9.8
AR-PFP	12.2 ± 5.7	
Workload		
Low	6.0 ± 1.0	32.2 ± 8.1
Moderate	8.0 ± 1.6	42.7 ± 10.8
High	9.4 ± 1.8	50.3 ± 12.3

SD, standard deviation; VO<sub>2PEAK</sub>, peak oxygen uptake; VO<sub>2PFP</sub>, average oxygen uptake for all tasks of modified CS-PFP; percent of VO<sub>2PEAK</sub>, VO<sub>2PFP</sub> as percentage of VO<sub>2PEAK</sub>; AR-PFP, VO<sub>2PEAK</sub> — VO<sub>2PFP</sub>; workload, average oxygen uptake for tasks of the low, moderate, and high workloads of the modified CS-PFP.

**Table 3.** Workload, task, and measurement categorisation of the CS-PFP\* and average  $\text{VO}_2^a$  during performance of the modified CS-PFP and values reported for specific tasks

Workload	Task	Measurement	$\text{VO}_2$ (ml/kg/min)	
			MOD CS-PFP	Reported data
Low	Weight carrying	Weight <sup>b</sup> , time	5.2 ± 1.2	
	Pouring	Time	5.9 ± 1.1	
	Shoe strapping	Time	5.6 ± 1.0	
	Scarves wrapping	Time	6.4 ± 1.2	
	Reaching	Distance	6.8 ± 1.4	
Moderate	Floor sweeping	Time	7.3 ± 1.6	8.8 <sup>c</sup> , 11.4 <sup>d</sup> , 11.7 <sup>g</sup>
	Laundry 1	Time	8.2 ± 1.8	7.0 <sup>c</sup>
	Laundry 2	Time	8.1 ± 1.7	
	Bed making	Time	10.0 ± 2.0	7.0 <sup>c</sup>
	Vacuuming	Time	8.0 ± 1.6	12.3 <sup>c</sup> , 9.8 <sup>d</sup> , 10.4 <sup>g</sup> , 8.1–8.9 <sup>e</sup>
	Floor down/up	Time	8.2 ± 1.4	
	Fire door	Time	6.8 ± 1.6	
High	Bus	Weight <sup>b</sup> , time	9.2 ± 1.7	
	Groceries	weight <sup>b</sup> , time	10.8 ± 2.4	8.8 <sup>c</sup> , 11.4 <sup>d</sup> , 11.7 <sup>g</sup>
	Stair climbing	Time	9.6 ± 1.8	
	Usual pace walking (3 min)	Distance	8.3 ± 2.4	8.8 <sup>f</sup>

\* CS-PFP, continuous-scale physical functional performance test.

<sup>a</sup>  $\text{VO}_2$  values include performance of the task and the time between tasks (generally standing, listening to instruction).

<sup>b</sup> CS-PFP<sub>MOD</sub>, modified CS-PFP (weight standardised for each task using weight in the CS-PFP; weight carrying, 14.75 lbs; bus, 12 lbs; and groceries, 14 lbs), reported data from <sup>c</sup> [18, 19], <sup>d</sup> [20], <sup>e</sup> [21], <sup>f</sup> [17], and <sup>g</sup> [22]. Values for this study are reported as means ± standard deviation and means from reported literature are listed.

## Discussion

The principle finding of this study was that higher aerobic capacity provided a greater aerobic reserve, which in turn, was associated with higher physical functional performance capacity in older adults. A higher aerobic reserve provides older adults with a larger margin of safety [4], which may ultimately attenuate functional decline, preserving independence.

A higher aerobic reserve allowed the older adults in this study to use a lower percentage of  $\text{VO}_{2\text{PEAK}}$  for functional performance (Table 2). The percentage of oxygen uptake elicited during the moderate and high workload levels (Table 2) are above the 40% intensity reported [6] as a sustainable workload throughout the day. Each workload, moderate and high, reflects an average oxygen uptake increase of 7.6–10.5% of  $\text{VO}_{2\text{PEAK}}$  (Table 2). A higher aerobic reserve allows older adults to do more throughout the day without encroaching upon their maximal aerobic capacity. These findings can be used to understand how changes in living status could affect the functional performance of older adults. For example, typical services provided by retirement communities range from weekly linen services to light housekeeping, eliminating the need to perform the moderate and high workload tasks included in the CS-PFP. Reducing the number of tasks that require strength can exacerbate declines in muscle mass and strength [3]. Without light housekeeping services the daily energy demand is lowered between 10 and 18% (from Table 2). Since the low workload tasks, on average, required only 6 ml/kg/min, or 32% of  $\text{VO}_{2\text{PEAK}}$ , older adults living in these facilities

should be aware of the need to engage in moderate intensity activities to maintain minimum recommended aerobic activity levels. Moderate-intensity walking and resistance training programmes have led to improvements in  $\text{VO}_{2\text{PEAK}}$  [23] and physical function [8, 23–27]. Maintaining an aerobic reserve may decrease functional limitations and delay the onset of dependence. In older adults, a positive correlation exists between aerobic capacity and daily physical activity [28]. Thus, older adults with a large aerobic reserve are more likely to have discretionary energy, which may allow them to engage in activities of interest and perform tasks necessary to maintain independence.

Recently published guidelines for physical activity for older adults recommend a systematic, well-rounded exercise programme in addition to routine activities of daily living that may include chores of light or moderate intensity [27]. Data presented here provide evidence that light household tasks and carrying groceries require moderate (40–60%) [13] aerobic intensity. The normal day consists of linking together multiple tasks and performing them consecutively while incorporating small intermittent breaks [6]. A systematic and scientific exercise programme where the heart rate is continuously elevated for thirty or more minutes a day could result in a significant increase in  $\text{VO}_{2\text{PEAK}}$ , and therefore, better aerobic reserve. In this way, the aerobic cost of household chores becomes a lower percentage of the overall aerobic capacity, and therefore, requires less effort.

Previously reported oxygen uptake values for performance of daily tasks are similar to those reported for the CS-PFP<sub>MOD</sub> (Table 3, marked) [18–22]. Average  $\text{VO}_2$  of the five values for the tasks labelled ‘c’ in Table 3 ( $8.8 \pm 2.2$

ml/kg/min) appears to be similar to the average  $\text{VO}_2$  for comparable tasks in the CS-PFP<sub>MOD</sub> ( $8.9 \pm 1.5$  ml/kg/min). Average oxygen uptake for tasks in the moderate workload ( $8.0 \pm 1.6$  ml/kg/min, Table 3) are slightly lower than the average for similar tasks reported in the literature ( $9.7 \pm 0.7$  ml/kg/min, Table 3), with high workload tasks ( $9.4 \pm 1.8$  ml/kg/min, Table 3) also slightly lower than tasks reported in the literature (Table 3) ( $11.2 \pm 2.4$  ml/kg/min). Differences in  $\text{VO}_2$  values for individual tasks may be attributed to differences in task execution and administration (e.g. muscle group used; vacuum model). The CS-PFP<sub>MOD</sub> tasks are administered to reflect how tasks are performed in daily life, short, non-steady state activities. Rest periods between tasks of the CS-PFP<sub>MOD</sub> most likely decreased the average  $\text{VO}_2$  value. However, the tasks labelled 'c' in Table 3, which had a higher average  $\text{VO}_2$  value, were performed for three or more minutes in order to achieve steady state oxygen uptake. Understanding the aerobic cost of serially performed tasks can provide researchers with insight into how much aerobic fitness is needed to optimise the living arrangements for older adults as aerobic capacity wanes in the later years of life.

While the mechanisms for age-related decreases in aerobic capacity are not fully understood, evidence from animal [29] and human [30–32] research supported that muscle taken from older research subjects had the capacity to make large energetic changes in response to both endurance and resistance training. Peripheral oxygen carrying capacity is improved with increases in the capillary-to-fibre ratio [30, 32] and increased blood flow to the working muscle at both maximal and sub-maximal capacities [33] may account for the improved performance in those with adequate aerobic reserve.

Some of the limitations of this study include the mask of the Cosmed K4b<sup>2</sup> unit that may interfere with vision during tasks such as sweeping the floor or climbing the stairs, which can lengthen the time taken to complete the task. The ability to generalise this research is limited to the mostly Caucasian female sample used in this study. However, it should be noted that the CS-PFP scores of this group are comparable to scores from previous research samples [34, 35]. Future investigations could implement training interventions to determine the effect of increased physical activity on aerobic reserve, functional performance and discretionary energy in older adults.

## Acknowledgements

The authors would like to thank Tanya Misko, Ed.D. and Jill Slade, Ph.D. for their help in the completion of this project.

## Key points

- In older adults, performance of daily tasks requires between 30 and 50% of aerobic capacity.
- Maintaining an aerobic capacity of 20 ml/kg/min allows for moderate and high workload tasks to be performed with 50% of aerobic capacity in reserve.

## Aerobic reserve and functional performance

- Higher aerobic reserves may provide more discretionary energy for work and leisure over and above that required to perform household tasks.

## Funding

This study was supported by a Georgia Gerontology Consortium Seed Grant.

## Conflict of interests

The authors of this article have no conflicts of interest in sources of funding or provenance of laboratory or biochemical equipment.

The data presented in this article have been published as an abstract.

## References

1. Cress ME, Meyer M. Maximal voluntary and functional performance levels needed for independence in adults aged 65 to 97 years. *Phys Ther* 2003; 83: 37–48.
2. Morey MC, Pieper CF, Cornoni-Huntley J. Physical fitness and functional limitations in community-dwelling older adults. *Med Sci Sports Exerc* 1998; 30: 715–23.
3. Posner JD, McCully KK, Landsberg LA, *et al.* Physical determinants of independence in mature women. *Arch Phys Med Rehabil* 1995; 76: 373–80.
4. Young A. Exercise physiology in geriatric practice. *Acta Med Scand Suppl* 1986; 711: 227–32.
5. Schwartz RS. Sarcopenia and physical performance in old age: introduction. *Muscle Nerve Suppl* 1997; 5: S10–2.
6. Astrand P, Rodahl K. *Textbook of Work Physiology: Physiological Bases of Exercise*, 2nd edition. New York: McGraw-Hill, 1977.
7. Cress ME, Buchner DM, Questad KA, *et al.* Continuous-scale physical functional performance in healthy older adults: a validation study. *Arch Phys Med Rehabil* 1996; 77: 1243–50.
8. Cress ME, Buchner DM, Questad KA, *et al.* Exercise: effects on physical functional performance in independent older adults. *J Gerontol A Biol Sci Med Sci* 1999; 54: M242–8.
9. Binder EF, Birge SJ, Spina R, *et al.* Peak aerobic power is an important component of physical performance in older women. *J Gerontol A Biol Sci Med Sci* 1999; 54: M353–6.
10. Hausswirth C, Bigard AX, Le Chevalier JM. The Cosmed K4 telemetry system as an accurate device for oxygen uptake measurements during exercise. *Int J Sports Med* 1997; 18: 449–53.
11. McLaughlin JE, King GA, Howley ET, *et al.* Validation of the COSMED K4 b2 portable metabolic system. *Int J Sports Med* 2001; 22: 280–4.
12. Cosmed S.R.L. K4b<sup>2</sup> User Manual. Rome, Italy: Cosmed S.R.L., 1998; 47–58.
13. American College of Sports Medicine, ACSM. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*, 4th edition. Baltimore, MD: Lippincott Williams and Wilkins, 2001.
14. Tanaka H, Desouza CA, Jones PP, *et al.* Greater rate of decline in maximal aerobic capacity with age in physically active vs. sedentary healthy women. *J Appl Physiol* 1997; 83: 1947–53.

**S. W. Arnett et al.**

15. Meredith CN, Frontera WR, Fisher EC, *et al.* Peripheral effects of endurance training in young and old subjects. *J Appl Physiol* 1989; 66: 2844–9.
16. Borg G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med* 1970; 2: 92–8.
17. Bard G, Ralston HJ. Measurement of energy expenditure during ambulation, with special reference to evaluation of assistive devices. *Arch Phys Med Rehabil* 1959; 40: 415–20.
18. Ainsworth BE, Haskell WL, Leon AS, *et al.* Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993; 25: 71–80.
19. Ainsworth BE, Haskell WL, Whitt MC, *et al.* Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000; 32: S498–504.
20. Gunn SM, Brooks AG, Withers RT, *et al.* Determining energy expenditure during some household and garden tasks. *Med Sci Sports Exerc* 2002; 34: 895–902.
21. Norman JF, Kautz JA, Wengler HD, *et al.* Physical demands of vacuuming in women using different models of vacuum cleaners. *Med Sci Sports Exerc* 2003; 35: 364–9.
22. Gunn SM, Brooks AG, Withers RT, *et al.* The energy cost of household and garden activities in 55- to 65-year-old males. *Eur J Appl Physiol* 2005; 94: 476–86.
23. Moore TL, Speer E, Johnson F, *et al.* The effects of aerobic training and nutrition education on functional performance in low socioeconomic older adults. *J Geriatric Phys Ther*, In press.
24. Miszko TA, Cress ME, Slade JM, *et al.* Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 2003; 58: 171–5.
25. Ades PA, Savage P, Cress ME, *et al.* Resistance training improves performance of specific measured daily activities in disabled women with coronary heart disease. *Med Sci Sports Exerc* 2003; 35: 1265–70.
26. Haykowsky M, McGavock J, Vonder Muhll I, *et al.* Effect of exercise training on peak aerobic power, left ventricular morphology, and muscle strength in healthy older women. *J Gerontol A Biol Sci Med Sci* 2005; 60: 307–11.
27. Cress ME, Buchner DM, Prohaska T, *et al.* Best practices for physical activity programs and behavior counseling in older adult populations. *J Aging Phys Act* 2005; 13: 61–74.
28. Starling RD, Toth MJ, Carpenter WH, *et al.* Energy requirements and physical activity in free-living older women and men: a doubly labeled water study. *J Appl Physiol* 1998; 85: 1063–9.
29. Kerner J, Turkaly PJ, Minkler PE, *et al.* Aging skeletal muscle mitochondria in the rat: decreased uncoupling protein-3 content. *Am J Physiol Endocrinol Metab* 2001; 281: E1054–62.
30. Hepple RT, Mackinnon SLM, Goodman JM, *et al.* Resistance and aerobic training in older men: effects on VO<sub>2</sub>peak and the capillary supply to skeletal muscle. *J Appl Physiol* 1997; 82: 1305–10.
31. Haykowsky M, McGavock J, Vonder Muhll I, *et al.* Muscle strength in healthy older women: Effect of exercise training on peak aerobic power, left ventricular morphology, and. *J Gerontol A Biol Sci Med Sci* 2005; 60: 307–11.
32. Jubrias SA, Esselman PC, Price LB, *et al.* Large energetic adaptations of elderly muscle to resistance and endurance training. *J Appl Physiol* 2001; 90: 1663–70.
33. Beere PA, Russell SD, Morey MC, *et al.* Aerobic exercise training can reverse age-related peripheral circulatory changes in healthy older men. *Circulation* 1999; 100: 1085–94.
34. Petrella JK, Miller LS, Cress ME. Leg extensor power, cognition, and functional performance in independent and marginally dependent older adults. *Age Ageing* 2004; 33: 342–8.
35. Petrella JK, Cress ME. Daily ambulation activity and task performance in community-dwelling older adults aged 63–71 years with preclinical disability. *J Gerontol A Biol Sci Med Sci* 2004; 59: 264–7.

Received 10 April 2007; accepted in revised form 30 November 2007